

# Ring Leader™ 17% + P Meal

## MEDICATED

A complete feed for pigs weighing 150 lbs.

For increased rate of weight gain, improved feed efficiency and increased carcass leanness in finishing swine, weighing not less than 150lbs, fed a complete ration containing at least 16% crude protein for the last 45 to 90 lb. of gain prior to slaughter.

Prevents the development of house flies, stable flies, face flies, and horn flies in the manure of treated swine.

### ACTIVE DRUG INGREDIENT

Ractopamine hydrochloride ..... 9 g/ton  
Diflubenzuron ..... 5 g/ton

### Guaranteed Analysis

Crude Protein		(Min)	17.0 %
Lysine		(Min)	1.10 %
Crude Fat		(Min)	5.4 %
Crude Fiber		(Max)	4.0 %
Calcium (Ca)	(Min) 0.7	(Max)	1.2 %
Phosphorus (P)		(Min)	0.71 %
Salt (NaCl)	(Min) 0.3	(Max)	0.8 %
Sodium	(Min) 0.1	(Max)	0.6 %
Selenium (Se)		(Min)	0.4 PPM
Zinc (Zn)		(Min)	986 PPM

### Feeding Directions

Feed as sole ration to pigs weighing 150 lbs.

### Ingredients

Grain Products, Plant Protein Products, Processed Grain By-Products, Fish Meal, Calcium Carbonate, Dried Whey Product, Brewers Dried Yeast, Dicalcium Phosphate, Salt, L-Lysine, Zinc Oxide, L-Threonine, Propionic Acid (a preservative), Choline Chloride, Potassium Chloride, Bentonite, DL-Methionine, Copper Sulfate, Natural and Artificial Flavors, Calcium Carbonate, Diatomaceous Earth (flow agent), Hemicellulose Extract, Natural and Artificial Flavors, Mineral Oil, Magnesium Oxide, , Roughage Products, Dehydrated Pichia pastoris Fermentation Extract, Plant Protein Product, Yeast Extract, Hydrolyzed Yeast, Mineral Oil, Chromium Tripicolinate, Ferrous Sulfate, Zinc Sulfate, Zinc Amino Acid Complex, Manganese Amino Acid Complex, Vitamin E Supplement, Manganese Sulfate, Niacin Supplement, Selenium Yeast, d-Calcium Pantothenate, Copper Amino Acid Coomplex, Biotin, Riboflavin Supplement, Menadione Nicotinamides Bisulfite (Source of Vitamin K Activity), Vitamin D3 Supplement, Vitamin A Supplement, Vitamin B12 Supplement, Sodium Selenite, Pyridoxine Hydrochloride, Folic Acid, Thiamine Mononitrate, Ethylenediamine Dihydriodide, Dried Cheese, Lecithin, Calcium Silicate and Soybean Oil.

### Caution

CAUTION: Pigs fed Ractopamine Hydrochloride are at an increased risk for exhibiting the fatigued pig syndrome (also referred to as "slows", "subs" or "suspects"). Pig handling methods to reduce the incidence of downer pigs should be thoroughly evaluated prior to initiating use of Ractopamine Hydrochloride. Ractopamine may increase the number of injured and/or fatigued pigs during marketing. Not for animals intended for breeding.

LIMITATIONS: Feed continuously as the sole ration. Ractopamine Hydrochloride should be fed in a complete swine ration containing at least 16% crude protein from 150 lb. (68 kg) to 240 lb. (109 kg) bodyweight. No increased benefit has been shown when ractopamine concentrations in the diet are greater than 4.5 g/ton.

CAUTION: Keep out of reach of children. Discard empty container according to local regulations. Never reuse empty container.

This product contains added copper. Do not feed to sheep or other copper sensitive animals.

4140 ~PC 430920 4  
Net Weight 50 Lb (22.6 kg) Or Bulk  
Meal  
~BCFMT